

The treatment of MG

My Medication

Anticholinesterrase inhibitors

Mestinon

Ubretid

Efedrin

Other

Corticosteroids

Atsatioprin

Thymectomy: No

Yes date:

Other diseases / medications

Information about MG

MG is an autoimmune disease. The function of the neuromuscular junction is defected and due to this, the patient has a tendency to fatigue and increased weakness of voluntary muscles after their continued use. Rest will ease the symptoms.

Several drugs worsen the symptoms: morphin, tubocurare, penisillamin, kinin, benzodiatzepin, barbiturates, fentiatzin, klorokin, magnesium.

Myastenic crises: When the patient is unable to swallow tablets, give *neostigmine* 1–2 mg i.m.

Take Note! Koliergic crises (overdose of drugs) leads to muscle weakness.

*Helsinki 18.3. 2001 Kiti Müller, MD.
Ph.D., Expert of Neurology*

Responsible for my treatment

Hospital

Phone

Physician

Phone

Contact man

Name

Address

Phone



**-patient
information
card**

The owner of this card has (MG)
Myasthenia Gravis

Name

Identity number

Address

Phone

Finnish MG-Association
Suomen MG-yhdistys ry
BOX 187 00101 HELSINKI
FINLAND
<http://www.suomenmg-yhdistys.fi>
toimisto@suomenmg-yhdistys.fi